

# RELATIONSHIP BETWEEN QUERCETIN AND HEALTH FROM THE PERSPECTIVE OF EPIGENETICS

Nguyen Hoang Nam, Pham Minh Tri, Nguyen Thi Minh Nguyet✉

*Department of Nutrition and Food Science, Institute of Biotechnology  
and Food Technology, Industrial University of Ho Chi Minh City, Vietnam*

## ABSTRACT

**Background:** Research on free radicals has attracted scientific and medical attention for decades, especially regarding aging, disease, and infectious and non-infectious diseases. Exposure to free radicals from daily pollution and unhealthy lifestyles accelerates bodily damage and aging. These factors are primary contributors to gene and chromatin dysregulation, leading to chronic diseases such as metabolic disorders, allergies, cancer, etc.

**Scope and approach:** In the present literature, quercetin has been shown to have the ability to neutralize free radicals, reduce the formation of inflammatory cytokines, induce histamine release, and induce cell damage. It also inhibits the growth of some inflammatory bacteria in the body and reverses disease-related epigenetic changes. Furthermore, quercetin strengthens the immune system, promoting health and resilience against external threats. Its multifaceted role is discussed in light of recent epigenetic research.

**Key findings and conclusions:** Quercetin's therapeutic potential via epigenetic mechanisms is significant, but further research is needed to fully understand its ability to reverse pathological epigenetic changes.

**Keywords:** *epigenetic, quercetin, antioxidant, reactive oxygen species, autophagy*

-----

## I. INTRODUCTION

Free radicals are unstable molecules that can potentially damage body cells and tissues. They are generated from many causes like the mitochondrial electron transport chain, inflammatory reactions, enzymes (e.g., xanthine oxidase), drug metabolism, smoking, radiation exposure (X-rays, UV rays) [1]. They have been linked to the risk of osteoporosis and eye diseases including cataracts, glaucoma, keratoconus, dry eye, myopia, macular degeneration, and pterygium [2, 3]. Free radicals and allergic disease are associated with dysregulation of epigenetic mechanisms. Plant-based diets provide the body with compounds including EGCG, curcumin, quercetin, and genistein that reverse the epigenetic mechanism involving DNMT and HAT enzymes [4]. Here we are mainly interested in quercetin. In addition to being obtained from the diet, quercetin is also abundant in plants with traditional medicinal uses, such as *Physalis angulata* L., *Hibiscus sabdariffa* L., *Anoectochilus setaceus*, and *Psidium guajava* L.

✉ Corresponding author: Nguyen Thi Minh Nguyet  
Email: nguyenthiminhnguyet@iuh.edu.vn  
Doi: 10.56283/1859-0381/732.

Submitted: 6/7/2024    Revised: 23/8/2024  
Accepted: 6/11/2024  
Published online: 11/11/2024

These plants are known for treating diseases, offering anti-inflammatory, antibacterial, anti-toxic, anti-cancer, and immune system-regulating properties [5-8]. Quercetin promotes the body to form endogenous enzymes, which neutralize free radicals and detoxify by the Nrf2 signaling pathway [9]. Quercetin protects hepatocytes, preventing excessive accumulation of lipid droplets in the nonalcoholic fatty liver via the IRE1a/XBP1s pathway [10].

In the process of synthesizing documents and proofs, several research areas are also mentioned and focused, including:

- Epigenetics is the study of the mechanisms by which behavioural and

environmental factors can induce changes that affect gene regulation, including processes such as gene methylation, histone modification (acetylation), and micro RNAs (miRNAs) [11].

- Nutrigenomics and nutrigenetics study the effects of nutrition on gene regulation, the impact on individual health, and the risk of developing diseases [12]. In addition, nutrigenomics and nutrigenetics also help to understand better how components of the diet, directly or indirectly, affect metabolism and how gene regulation occurs in humans. More specifically, this research direction is also interested in effective ways to prevent diseases and reduce risks in disease treatment and medical costs [12].

## II. METHODS

We conducted a comprehensive literature review of quercetin, genetics, epigenetics, and the molecular relationship between diet and health. We searched Google Scholar, Springer, PubMed, and ScienceDirect using keywords such as epigenetics, quercetin, antioxidant, reactive oxygen species, and autophagy, focusing on publications from 2002 to 2024. From an initial pool of 352

articles, we selected 77 relevant to our research question. The remaining 275 were excluded due to language barriers (non-English or Vietnamese), irrelevance, paywalls, or questionable source reliability. We also searched for relevant reports in libraries of organizations such as the World Health Organization, the World Bank, research institutes, and universities.

## III. INTRODUCTION TO QUERCETIN

Quercetin, a compound found in various foods (Table 1), exists in multiple forms, including quercetin-3,4'-O-diglucoside, quercetin-3-O-glucoside, quercetin-3-O-rhamnoside, quercetin-4'-O-glucoside, quercetin-3-O-rutinoside [13].

Quercetin is also found in many plants that act as a medicine for many diseases including *Physalis angulata* L., *Hibiscus sabdariffa* L., *Anoectochilus setaceus*, *Psidium guajava* L., v.v [5-8].

*Physalis angulata* L. is distributed mainly in tropical and subtropical regions including Asia, Central and South America, Africa, and the Pacific Islands. *Physalis angulata* L. is edible and exhibits anti-inflammatory, antibacterial, anti-toxic, anti-cancer, and immune-modulating effects attributed to phenolic acids (chlorogenic acid, caffeic acid, p-coumaric acid) and flavonoids (rutin, quercitrin, quercetin and kaempferol) [6].

Hibiscus (*Hibiscus sabdariffa* L.) is used as food and medicine. Their calyxes are rich in flavonoids, especially quercetin and anthocyanin, with quercetin demonstrating antibacterial activity [5].

Guava (*Psidium guajava* L.) is distributed in tropical and subtropical regions such as Asia, Africa, South America, and the Caribbean. In Viet Nam, *Psidium guajava* L. is very popular and is grown in many places but mainly in the Mekong Delta with an area of nearly 4500 ha. Guava leaves, also used as a tea, contain quercetin-3-O-sulfate, which exhibit anti-diabetic, antioxidant,

anti-bacterial, anti-cancer, anti-diarrheal, and anti-inflammatory properties [8].

Fenugreek (*Trigonella foenum-graecum*), a herb widely grown in India, Pakistan, and the Middle East, possesses antibacterial, antiviral, anti-tumor, anti-inflammatory, and antioxidant effects. Both its leaves and seeds, widely consumed as food and medicine, are rich in nutrients like vitamins, iron,  $\beta$ -carotene, etc. Its seeds contain a variety of alkaloids, flavonoids, saponins, and carbohydrates. Leaf analysis revealed the presence of two main flavonoids, quercetin, and kaempferol [27].

**Table 1.** Quercetin content in food, beverages and health studies on quercetin.

Food and beverage	Quercetin	Research on quercetin for health	Ref.
Food (mg/100 g)			
Apple	4.01	Quercetin-3-glucoside (Q3G), coumaric acid, phloridzin, quercetin and phloretin are the major polyphenolic compounds that makeup apple pulp. Studies on HeLa cell lines show that Q3G inhibits the cell proliferation cycle in the S phase, promotes apoptosis by activating caspase 9/3, reduces the expression of BCL 2 anti-apoptosis protein, and upregulates the expression of BAX pro-apoptotic protein. The Q3G present in apple pulp holds promise as an anti-inflammatory and anticancer agent for the treatment of cervical cancer.	[13, 14]
Blueberry	5.05	The most abundant polyphenols detected in extracts of <i>V. myrtillus</i> were quercetin, kaempferol, phenolic acid, and gentisic acid. A study on the colorectal cancer cell line HCT-116 showed that quercetin and kaempferol have strong cytotoxic, antioxidant, and apoptotic effects.	[15, 16]
Garlic	1.74	Garlic oil contains 0.24 mg/g quercetin and 0.33 mg/g gallic acid. The study showed that garlic essential oil had antibacterial properties against <i>Staphylococcus aureus</i> ,	[17, 18]

Food and beverage	Quercetin	Research on quercetin for health	Ref.
		and the diameter of the growth inhibition zone increased from $20.20 \pm 0.58$ mm (37.5 mg/mL) to $31.70 \pm 0.55$ mm (300 mg/mL).	
Okra	20.97	Quercetin-3-gentiobioside (Q3G), quercetin-3-sambubioside (Q3S), rutin, quercetin-7-glucoside (Q7G), isoquercitrin (ISO), and quercetin-3-malonylglucoside (Q3M) are found in okra. The study showed that Q3G, Q3S, ISO, and Q3M significantly inhibited the proliferation of cell lines NCI-N87, A375, A549 (25- 100 $\mu$ mol/ L), and HFLS-RA (200- 300 $\mu$ mol/ L).	[17, 19]
Onions ( <i>Allium cepa</i> L.)	20.30	Neuronal ischemia is the cause of neuronal damage and gliosis. Research has shown that onion extract ( <i>Allium cepa</i> L.) and quercetin contribute to the amelioration of ischemic neuropathy, protecting hippocampal pyramidal cells in the CA1 region, simultaneously reducing 4-hydroxy-2-nonenal formation in a male Mongolian gerbil model ( <i>Meriones unguiculatus</i> ).	[17, 20]
Beverage			
Black tea	2.50 (mg/100 ml)	The study was based on venous blood samples from healthy adults (25-30 years old). Research has shown that BPA damages and causes swelling of red blood cells leading to hemolysis. Black tea extract and quercetin play a role in protecting red blood cells and significantly slowing down hemolysis caused by BPA.	[13, 21]
Red wine	3.16 (mg/ 100ml)	The skin cancer study on CD1 mice showed that red wine has 4 types of polyphenols including quercetin, gallic acid, (+)-catechin, and trans-resveratrol that play a role in cancer prevention. Research indicates that trans-resveratrol is absorbed much more efficiently than (+)-catechin and quercetin after oral administration, concluding that trans-resveratrol may be the most effective anti-cancer polyphenol present in red wine when taken orally consumed by healthy human subjects.	[13, 22]
Lotus leaf tea	105.93 (mg/ 100 g)	Overdose of acetaminophen resulted in mitochondrial damage leading to leakage of damaging free radicals in hepatocytes of AML (ATCC CRL-2254) mice.	[9, 23]

Food and beverage	Quercetin	Research on quercetin for health	Ref.
		The Nrf2 signaling pathway involved in the transcription of antioxidant and detoxifying enzymes including HO 1, NQO 1, and UGT 1A is activated by phenolic compounds (gallic acid, EGCG, myricitrin, and rutin), ellagic acid, myricetin, quercetin, luteolin, kaempferol and apigenin), compounds extracted from lotus leaves act as inhibitors of DNMT and HDAC enzymes.	
Pomegranate juice	1.456 (mg/100 g)	60 strains of <i>Staphylococcus epidermidis</i> isolated from human eye infections were obtained from the Instituto de Oftalmología Conde de Valenciana, Mexico City. Half of the strains were resistant to multiple drugs and other compounds, such as ethidium bromide and benzalkonium chloride.  Research has shown that pomegranate juice plays an antibacterial role against resistant bacteria <i>Staphylococcus epidermidis</i> .	[24, 25]
Prickly pears ( <i>Opuntia spp.</i> )	4.86 (mg/100 g)	The study was based on four cancer cell lines: mammary (MCF-7), prostate (PC3), colon (Caco2) and hepatic (HepG2).  Research shows that juice derived from Prickly pears ( <i>Opuntia spp.</i> ) plays a role in inhibiting the proliferation of cancer cells and fighting free radicals. <i>Opuntia violaceae</i> Moradillo contains the highest content of flavonoids and inhibits prostate and colon cancer cell proliferation.  <i>Opuntia rastrera</i> Rastrero inhibits the growth of four cancer cell lines.	[17, 26]

### 3.1. Quercetin inhibits cancer

Cancer, a leading global health concern, encompasses a group of related diseases characterized by abnormal cell growth and uncontrolled division. Unlike normal cells, cancerous cells do not die but proliferate, forming tumors and invading surrounding tissues. These cells can also metastasize through the circulatory or lymphatic systems. While cancer drugs

have been developed, a universal cure for all cancer types remains elusive [28].

Bahare Salehi et al. (2020) highlighted quercetin's therapeutic potential for diseases like diabetes, Alzheimer's, arthritis, parasitic infections, and cancer. Preclinical and clinical studies have explored its effects on various cancers, including pancreatic, osteosarcoma, ovarian, breast, cervical,


colon, and gastrointestinal cancers [29]. Sections 3.2 to 3.4 further discuss quercetin's potential role in cancer treatment, focusing on its bidirectional relationship with epigenetics. However, the application of quercetin in humans

faces challenges due to its poor water solubility, permeability, and instability in physiological environments, along with its short half-life and significant first-pass metabolism, leading to poor oral bioavailability [29].

### 3.2. Gastrointestinal cancer

Gastrointestinal cancer represented the largest proportion of new cancer cases worldwide in 2018 (26.3% of 4.8 million). Colorectal cancer was the most

common (10.2%), followed by stomach (5.7%), liver (4.7%), esophageal (3.2%), and pancreatic cancers (2.5%) (see Figure 1) [30].




**Figure 1.** Statistical data on the number of new cases of gastrointestinal cancer globally in 2018 out of a total of 4.8 million cases [30].

Regarding the mechanism of gastrointestinal tumour development, pathogens (such as viruses, bacteria, and parasites) have evolved over a long period, finding ways to efficiently enter the host and transform it into a host adaptive changes to increase their survival (e.g., *Helicobacter pylori* tolerating the acidic environment of the human stomach) [31]. Microorganisms host different pathologies such as exposure to *Helicobacter pylori* and Epstein-Barr Virus. *Helicobacter pylori* promote methylation of tumour suppressor genes including E-cadherin (*CDH1*), heart and neural crest

derivatives expressed 1 (*HAND1*), and multiple tumour suppressor 1 (p16 *INK4A*) [31]. Epstein-Barr virus promotes transcription and formation of DNA methyltransferase (DNMT) enzymes. These include DNMT 1, and DNMT 3A/3B formed by the transcription factor NF- $\kappa$ B. DNMTs involved in methylation of the tumour suppressor genes O-6-methylguanine-DNA methyltransferase (*MGMT*), Phosphatase and tensin homolog (*PTEN*) cause gastric cancer (see Figure 2) [32, 33]. In addition, smoking and alcohol consumption trigger inflammatory responses that lead to gastric damage,

increasing the risk of exposure to *Helicobacter pylori* and Epstein-Barr virus [34, 35].



**Figure 2.** Effect of *Helicobacter pylori* and Epstein-Barr virus on increased risk of stomach cancer.

Quercetin disrupts the cell barrier in gram-negative bacteria (e.g., *Escherichia coli*) via lipid peroxidation, causing cell lysis. Simultaneously, it activates the kynurenine pathway, depleting L-tryptophan and inhibiting growth of gram-positive bacteria [31]. Yali Liu et al. (2024) found that quercetin in liquorice exhibits anti-cancer properties by inhibiting proliferation, reducing tumor volume, inducing cell cycle arrest, and promoting apoptosis. These effects

correlate with increased Cyt-C, caspase-3, and BAX, decreased BCL-2 and Ki67, reduced mitochondrial membrane potential, and mitochondrial damage [36]. Therefore, quercetin may protect the stomach from inflammatory damage caused by external agents via its antibacterial activity. Furthermore, adopting a healthy lifestyle, limiting alcohol consumption, and avoiding smoking can further reduce the risk of digestive tract cancers.

### 3.3. Breast cancer

Breast cancer is a fairly common cancer in women worldwide with more than 2 million new cases in 2020 [37]. Breast

cancer is the fastest-growing cancer in India and has now surpassed cervical cancer. The National Cancer Registry

Program in 2018 estimated 162,468 new breast cancer cases and 87,090 deaths [38]. Breast cancer is known and documented in the "Edwin Smith Papyrus", about 3600 years ago (about 1600 BC). More recently, in the period of ancient Egyptian civilization, Hippocrates (460- 370 BC) suggested that the human body was an entity balanced by four types of humours including blood, phlegm, yellow bile, and black bile, representing respectively air, fire, earth, and water, help constitute all natural things. Breast cancer is the result of an imbalance, an excess of black bile, which causes tumors [39].

From the perspective of modern science and epigenetics, tumor suppressor genes play an important role in inhibiting or regulating cell proliferation by encoding proteins that block the action of tumor cell proteins. Tumor suppressor genes including *APC*, *BRCA1*, *Cyclin D2*, *GSTP1*, *p16*, *PTEN*, *RAR $\beta$* , *RASSF1A*, *ZMYND10* are among the important causes of breast tumors and are potential biomarkers in clinical applications, prognosis and early diagnosis of breast cancer [40].

Gene *BRCA1* is a tumour suppressor gene, which helps protect our body from uncontrolled proliferation of cells, leading to inhibiting the formation of cancers. Simultaneously, *BRCA1*, and *BRCA2* together with *MGMT* and *RAD 51* involved in the repair of the double-strand break of DNA [41]. Research shows that a diet low in folic acid and cobalamin increases the risk of tumour formation by increasing the rate of promoter methylation of the *RAR  $\beta$*  and *BRCA1* genes [42, 43].

### 3.4. Cervical cancer

Cervical cancer is a common malignancy in women annually worldwide with an

Research by Sai Kundur et al. (2018) has shown that quercetin and curcumin enhance histone acetylation at the H3K9 position of the promoter of the *BRCA1* gene [44]. Inhibition of cell proliferation and activation of apoptosis in the MDA-MB-231 (ATCC, HTB-26) cancer cell line was observed at a concentration of 20  $\mu$ M of quercetin. This medium and concentration also suppressed cell cycle progression through upregulation of the expression of FOXO3a by the JNK signaling pathway [45]. Tianyu Luo et al. (2024) proved that quercetin inhibited breast cancer progression in chronically stressed C57BL/6J mice by regulating microglia M1 polarization by targeting the STAT1 protein [46]. Marta et al. (2020) reported that quercetin inhibited Lef1 and reduced the expression of drug resistance-related proteins such as ABCG2, Vimentin, and Caveolin-1 on the docetaxel-resistant MCF7 breast cancer cell line. Quercetin was tested in a phase I clinical study in 1996, which demonstrated that it could be used safely in humans without significant toxicity at doses up to 1700 mg/m<sup>2</sup>. These findings support the feasibility of using quercetin as a safe adjuvant therapy in combination with docetaxel for the treatment of drug-resistant breast cancer, as it has the potential to enhance cancer cell sensitivity without posing significant risk to the patient [47].

From the above evidence, a quercetin-rich diet plays a very important role in breast cancer suppression through inhibition of cell proliferation, activation of apoptosis, and alteration of epigenetic mechanisms for tumor suppressor genes.

estimated 493,000 cases and more than 273,500 deaths. Methylation of the



promoter of the tumour suppressor gene *RASSF1A* is associated with a variety of cancers in breast, liver, pancreas, prostate, kidney, brain, stomach, blood, skin, lymphatic system (e.g., Hodgkin's lymphoma), etc. Compounds methionine, vitamin B<sub>12</sub>, resveratrol, curcumin, emodin, peperomin E, dioscin, mahanine, and PEITC participate in the demethylation of the promoter of the *RASSF1A* gene [48]. Studies on HeLa cell lines show that quercetin and eugenol enhance the demethylation of the *RASSF1A* gene [49].

In addition, FADD and caspase 8 participate in apoptosis, inhibiting cervical cancer formation. HPV-E6 virus inhibits apoptosis through HPV-E6 binding and degradation of FADD

leading to an increased risk of cervical cancer [50]. Flavonoid compounds (myricetin, morin, quercetin, kaempferol, galangin, etc.) act as inhibitors of the interaction between HPV-E6 and FADD, caspase 8 [51]. Motoki Murata et al. (2022) admitted that dietary quercetin enhanced the expression of cervical tumour-inhibiting miRNAs, including miR-26b, miR-126, and miR-320a, and simultaneously inhibiting the *CTNNB1* gene, which encodes  $\beta$ -catenin, by upregulating miR-320a expression [52].


From the above evidence, it is shown that a diet rich in quercetin also plays a strategic role in inhibiting cervical cancer by activating apoptosis and altering the epigenetic mechanism of tumor suppressor genes.

#### IV. QUERCETIN FOR TYPE 2 DIABETES AND AGE

Advanced glycation end products (AGEs) are a complex group of compounds formed in multiple stages with the Maillard reaction playing a key role. AGE is generated at the last stage of the Maillard reaction through polymerization, oxidation, dehydration, and cyclization events, discovered by Louis-Camille Maillard [53].

Type 2 diabetes is recognized as a global epidemic in modern times, causing a heavy burden of public health concerns worldwide, forecasting 783.2 million cases in adults aged 20–79 years in 2045 [54]. High blood sugar is responsible for

endogenous AGE synthesis. AGE together with the RAGE receptor activates inflammatory responses (NF- $\kappa$ B, NOX1, ERKs, Janus kinases, MAPK) that damage cells leading to diseases including retinopathy, neuropathy, nephropathy, and atherosclerosis. Endogenous AGE results from the high blood glucose level in patients with diabetes, while exogenous AGE originates from a typical Western diet with the characteristics of high-temperature food processing and cooking for a long time, low hydration, and high pH [53].



**Figure 3.** Relationship between quercetin, and exercise for HDAC5 in promoting the enhancement of GLUT4 expression through histone acetylation in disease prevention.


Exercise and a quercetin-rich diet have a major effect on phosphorylation at the Thr 172 site of AMPK. AMPK again activates phosphorylation at Ser 259 and Ser 498, leading to inhibition of HDAC5 enzyme activity. This development contributes to promoting histone acetylation, helping to untwist the chromatin of the *GLUT4* gene in skeletal muscle cells, contributing to reducing

blood sugar levels, and improving the condition in diabetics [55, 56]. Nuclear factor erythroid 2-related factor 2 (Nrf2), a key transcription factor, plays a crucial role in regulating antioxidant and anti-inflammatory responses, thereby protecting cells from oxidative damage and inflammation-induced injury. Quercetin ameliorated diabetic kidney disease by inhibiting ferroptosis and

modulating Nrf2 in streptozotocin-induced diabetic rats [57]. AGE has been reported to inhibit osteoblastogenesis and bone mineralization, increasing the risk of osteoporosis by promoting the expression of DNA methyltransferase enzyme (DNMT1/3A/3B) leading to methylation of promoter regions in genes like alkaline phosphatase (*ALP*), osteopontin (*OPN*), osteocalcin (*OCN*), and Runt-related transcription factor 2 (*RUNX2*) (see Figure 3) [58].

From the above evidence, inflammatory reactions triggered by endogenous AGEs are linked to complications in patients with diabetes. In addition, limiting Western-style diets, reducing processing, and heating foods to high temperatures contribute to a reduction in exogenous AGE formation. Exercise and a diet rich in quercetin contribute to the activation of AMPK, which inhibits the HDAC5 enzyme leading to a reversal of the epigenetic mechanism that improves and reduces the progression of diabetes.

## V. THE ROLE OF QUERCETIN IN AUTOPHAGY RELATED TO NAFLD AND TUBERCULOSIS



**Figure 4.** Role of a diet rich in sulforaphane and quercetin-activating autophagy in the prevention of NAFLD.

Autophagy is a cellular process that degrades and recycles intracellular components like damaged organelles, misfolded protein molecules, lipid droplets, and killed pathogens [59]. Rajat Pant et al. (2023) demonstrated that a

high-fat and high-fructose diet in mice significantly increased the number of Kupffer cells with an M1-dominant phenotype in mice leading to the activation of inflammatory responses. This diet also increased the expression of

the DNMT1 enzyme, leading to methylation of LC 3B, ATG 5, and ATG 7 genes, which disrupted the degradation of lipid droplets (lipofautophagy), resulting in excessive lipid accumulation in hepatocytes. This is the basic cause of non-alcoholic fatty liver disease (NAFLD) [60].

Conversely, dietary sulforaphane, which activates the Nrf2 signaling pathway, promotes autophagy gene transcription (*LC3*, *ATG5*, and *ATG7*) in hepatocytes [61]. A diet rich in quercetin suppresses the progression of atherosclerosis by enhancing autophagy activity in the foam cells of RAW 264.7 macrophages [62], preventing excessive lipid droplet accumulation in the nonalcoholic fatty liver via the IRE1 $\alpha$ /XBP1s pathway (see Figure 4) [10]. Thus, while a high-fat, high-fructose diet disrupts autophagy through DNMT1, contributing to NAFLD, sulforaphane and quercetin enhance autophagy via Nrf2 and IRE1 $\alpha$ /XBP1s, respectively.

*Mycobacterium tuberculosis*, a causative agent of tuberculosis, causes 10 million new cases and 1.2 million deaths each year. Patients will develop symptoms including a persistent cough and can spread the disease through airborne particles containing *M. tuberculosis* [63]. Tuberculosis has been reported to increase its ability to survive and spread to the host through epigenetic and hibernating mechanisms [63, 64].


In an epigenetic mechanism, *Mycobacterium tuberculosis* inhibits and disrupts autophagy by enhancing the activity of G9a, EZH2, and HDAC3 enzymes. G9a and EZH2 mediate

respectively methylation of the histone tail H3K9 (H3K9 me2/3) and H3K27 (H3K27 me3), and they reduce the acetylation in the histone tails H3K9 (H3K9 ac) and H3K27 (H3K27 ac) by HDAC3. These processes lead to chromatin closure, transcriptional inhibition, and reduced expression of ATG 5 and ATG 7 genes involved in autophagy in RAW 264.7 macrophages (see Figure 5) [64].

Under favorable environmental conditions, with adequate nutrients, glucose participates in glycolysis and the TCA cycle leading to the breakdown of glucose into ATP, providing energy for the life of *M. tuberculosis* [63].

Under nutrient-deficient conditions, the isocitrate lyase converts isocitrate to glyoxylate in the glyoxylate cycle, facilitating lipid anabolism via the gluconeogenesis pathway. Here lipid acts as an energy source for hibernation. *M. tuberculosis*'s metabolic adaptability enables antibiotic resistance, facilitating transmission and survival within hosts, resulting in lengthy treatment and high reinfection rates [63]. An important finding was that isocitrate lyase inhibition by quercetin bound to the N-terminus resulted in the inhibition of Tuberculosis bacteria growth at concentrations of 3.57  $\mu$ M(IC<sub>50</sub>) [65].

In summary, autophagy plays a crucial role in host defense against disease. *M. tuberculosis* circumvents these defenses via epigenetic dysregulation and hibernation. Quercetin inhibits *M. tuberculosis* survival during hibernation by targeting isocitrate lyase.



**Figure 5.** *Mycobacterium tuberculosis* enhances survival by disrupting autophagy and hibernation, with quercetin playing a protective role against infectious disease.

## VI. QUERCETIN PROTECTS THE BODY FROM RADICALS


Free radicals are atoms, molecules, or ions with unpaired electrons that are highly active in chemical reactions with other molecules. In biological systems, free radicals are usually derived from oxygen (ROS), nitrogen (RNS), and sulfur (RSS) [1]. Reactive oxygen species (ROS) are formed from many sources including mitochondrial electron transport chains, inflammatory responses, enzymes such as xanthine oxidase, drug metabolism, smoking, exposure to X-rays and UV rays, etc. [1]. According to Liangping Li et al. (2021), ROS indirectly enhances the activity of the DNA-methyltransferase 3 beta enzyme leading to methylation of the promoter of the Kruppel-like factor 5 (*KLF5*) gene, inhibiting the wingless/int-1 (Wnt) signalling pathway, reducing the ability to differentiate osteoblasts leading to an increased risk of osteoporosis [3]. ROS damages the eye organs including the lens, cornea, anterior chamber, retina, ocular surface, optic nerve, and macula leading to eye diseases including

cataracts, glaucoma, keratoconus, dry eye, myopia, macular degeneration, and pterygium. Ultraviolet rays (UV), light-emitting diodes (LED), and electronic devices increase eye diseases [2]. Myopia is a common eye disease affecting approximately 1.6 billion people worldwide in 2000, GSTP1 and TXNRD2 enzymes participate in the neutralization of free radicals, contributing to the protection of the lens epithelium to help protect the lens from the risk of cataracts in highly myopic eyes. ROS promotes methylation of the GSTP1 and TXNRD2 promoter genes by the enzyme DNMT1, leading to reduced formation of free radical-neutralizing enzymes [66, 67]. A plant-based diet consisting of ascorbic acid,  $\alpha$ -tocopherol,  $\beta$ -carotene, lutein, zeaxanthin, and quercetin protects the eyes from free radicals [2, 68].

According to Zheng-Yuan Su et al. (2022), paracetamol (acetaminophen, APAP) is a common pain and fever reliever, their use in overdose damages

mitochondria leading to leakage of free radicals that cause hepatocyte damage. The Nrf2 signaling pathway involved in the transcription of antioxidant and detoxifying enzymes including HO 1, NQO 1, and UGT 1A is activated by phenolic compounds (gallic acid, EGCG,

myricitrin, rutin, ellagic acid, myricetin, quercetin, luteolin, kaempferol, and apigenin). Compounds extracted from lotus leaves act as inhibitors of DNMT and HDAC enzyme activity (see Figure 6) [9].



**Figure 6.** The role of a phenol-rich diet that inhibits HDAC and DNMT enzymes in protecting the body from ROS in the prevention of diseases.

From the above evidence, free radicals are the main cause of epigenetic disorders, eye diseases, and liver damage caused by paracetamol overdose. Diets rich in phenols including gallic acid, EGCG, myricitrin, rutin, ellagic acid, myricetin, quercetin, luteolin,

kaempferol, and apigenin plant-derived contribute to Nrf2 activation involved in the transcription of endogenous enzymes (antioxidant and detoxifying) through inhibition of HDAC and DNMT helps to protect eyes, liver, and other organs in the body from free radical damage.

## VII. QUERCETIN PROTECTS NEURONS FROM TRAUMATIC BRAIN INJURY


Traumatic brain injury affects more than 10 million people worldwide each year. Traumatic brain injuries caused by external forces (traffic accidents, falls, attacks) damage the brain and the blood-brain barrier. Trauma causes disturbances of a multitude of biochemical and

metabolic functions that affect brain cell homeostasis and lead to temporary or permanent impairment of consciousness, neuromotor disabilities, and psychological disturbances [69].

The study noted traumatic brain injury disrupted the epigenetic mechanism and

enhanced histone acetylation of the *HMGB1* gene by the enzyme HAT. HMGB1 binds to microglia's Toll-like receptors (TLRs) leading to activation of the inflammatory response (NF- $\kappa$ B) signalling pathway, which damages nerve cells and worsens the disease [70, 71]. A

diet rich in omega 3 activates the SIRT 1 enzyme, SIRT 1 enhances the expression of HDAC enzyme involved in histone deacetylation of the *HMGB1* gene and HMGB1 protein, which contributes to inhibiting the inflammatory pathway of HMGB1/ NF - $\kappa$ B (see Figure 7) [70, 72].



**Figure 7.** The role of omega 3 and quercetin in protecting the brain from traumatic brain injury.

Traumatic brain injury damages mitochondria, which leak free radicals that lead to the activation of inflammatory responses that damage nerve cells. Compounds including glutathione, N-acetylcysteine, ascorbic acid, flavonoids, coenzyme Q10, carotenoids, resveratrol, omega 3,  $\alpha$ -tocopherol, and more help neutralize free radicals [69]. Quercetin plays an anti-apoptotic role, neutralizing free radicals through activation of

Glutathione Peroxidase (GSH-Px), Superoxide Dismutase (SOD), and Catalase (CAT) enzymes and reducing inflammatory cytokines caused by traumatic brain injury [73]. Hitler Kom et al. (2019) admitted that quercetin treatment reduced the level of lipid peroxidation and increased SOD, CAT, and GSH-Px activities in the cortical area compared to the inflamed group. These findings suggest that dietary quercetin may help mitigate neuroinflammation-

induced oxidative stress as well as associated behavioural and histological changes [74].

From the above evidence, it is shown that traumatic brain injury disrupts the epigenetic mechanism, enhances histone acetylation of the *HMGB1* gene by the HAT enzyme, and promotes the activation of the inflammatory pathway

of HMGB1/NF- $\kappa$ B. A diet rich in glutathione, ascorbic acid, flavonoids, coenzyme Q10, carotenoids, resveratrol, omega 3,  $\alpha$ -tocopherol, quercetin, etc contributes to the neutralization of free radicals caused by mitochondrial damage. Diets rich in omega 3 play a role in reducing brain damage by reversing epigenetic mechanisms.

## VIII. QUERCETIN PROTECTS THE BODY FROM ALLERGIES

Allergic diseases include allergic rhinitis, asthma, food allergies, drug allergies, atopic dermatitis and urticaria, and angioedema. These different allergic diseases have in common associated with an individual's hypersensitivity to foreign agents leading to overactivity of the immune system. Immune system overactivity is associated with epigenetic dysregulation of the DNMT1 enzyme with an emerging role in allergic diseases [75].

### 8.1 Allergic rhinitis

Quercetin is a bioactive compound implicated in reducing the progression of human diseases, such as malignancies, metabolic syndrome, and autoimmune, and allergic diseases. In particular, allergic rhinitis is a common inflammatory condition that affects a large number of people around the world. The mechanism of allergic rhinitis is quite complex. Th2 cells are IL-4 and IL-5-releasing agents, which facilitate the release of allergenic, IgE by Mast cells. In

Polyphenol compounds (EGCG, curcumin, quercetin, genistein) reverse the epigenetic mechanism involving DNMT and HAT enzymes of the immune system. EGCG inhibits DNMT1 and HDAC enzymes, curcumin inhibits the HAT enzyme, quercetin activates the HAT enzyme and inhibits the HDAC enzyme, and genistein activates the HAT enzyme and inhibits the SIRT enzyme [4].

addition, Th17 cells promote inflammatory responses, while Treg regulates anti-inflammatory responses. The imbalance between Th1/Th2 and Treg/Th17 is the cause leading to allergic rhinitis. Quercetin plays a role in balancing the immune system by reducing Ig E, Ig G1, and histamine and increasing Ig G2 in serum, enhancing the expression and activity of FOXP3 and IL 10 which contributes to the reduction of inflammation [76].

### 8.2 Food allergies

Food allergy is an immune-mediated adverse reaction involving two classical mechanisms, IgE-mediated and non-IgE-mediated responses to symptoms involving the skin, gastrointestinal tract, respiratory tract, and cardiovascular


system. Almost any food can cause an allergic reaction, but 90% of recorded food allergies are caused by 8 food groups including eggs, milk, fish, shellfish, peanuts, soybeans, tree nuts, and wheat [77].



Th1/Th2 epigenetic dysregulation has been implicated in neonates. After birth, Th2 predominance is attenuated and Th1 is gradually increased, accompanied by demethylation leading to enhanced IFN $\gamma$  and IL10 expression for Treg. Abnormal changes that disrupt epigenetic regulation leading to disruption of the normal development of the immune system are involved in the phenotype of allergic disease (see Figure 8) [77].

From the above evidence, it is shown that epigenetic dysregulation leads to

disruption of the normal development of the immune system related to the phenotype of allergic disease. Plant-based diets rich in polyphenols (EGCG, curcumin, quercetin, genistein) have an impact on DNMT and HAT enzymes that reverse the epigenetic mechanism of the immune system, contributing to the immune system balance between Th1/Th2 and Treg/Th17 resulted in decreased inflammation and decreased IL 4, IL 5, Ig E and histamine.



**Figure 8.** Epigenetic dysregulation of IFN $\gamma$ , IL10, IL4/5, and FOXP3 genes related to the balance of Th1/Th2 related to the phenotype of allergic disease [77].

## IX. CONCLUSION

A diet rich in fruits, vegetables, seeds, beans, whole grains, and juices contains high levels of biological compounds including quercetin that may prevent cancer, minimize the formation of harmful free radicals in the body, prevent eye diseases, osteoporosis, cancer, anti-inflammatory, anti-allergic (histamine, IL 4/5, IgE), inhibiting gram (+), gram (-) bacteria, reversing epigenetic mechanisms of the disease, enhanced

autophagy to prevent nonalcoholic fatty liver disease.

A plant-rich diet not only benefits human health but also contributes to environmental protection, saving resources, reducing pollution, and conserving biodiversity. Therefore, we should give priority to eating foods from nature, limiting foods of animal origin, industrial foods, high-temperature processing, and increasing exercise to improve quality.

## References

1. Nega M. Role of Endogenous and Exogenous Glutathione in the Detoxification of Free Radicals. *Research & Reviews: A Journal of Toxicology*. 2017;7(1):1- 14.
2. Hsueh YJ, Chen YN, Tsao YT, Cheng CM, Wu WC, Chen HC. The Pathomechanism, Antioxidant Biomarkers, and Treatment of Oxidative Stress-Related Eye Diseases. *International journal of molecular sciences*. 2022;23(3). doi:10.3390/ijms23031255.
3. Li L, Wang H, Chen X, Li X, Wang G, Jie Z, et al. Oxidative Stress-Induced Hypermethylation of KLF5 Promoter Mediated by DNMT3B Impairs Osteogenesis by Diminishing the Interaction with  $\beta$ -Catenin. *Antioxid Redox Signal*. 2021;35(1):1-20. doi:10.1089/ars.2020.8200.
4. Cuevas A, Saavedra N, Salazar LA, Abdalla DS. Modulation of immune function by polyphenols: possible contribution of epigenetic factors. *Nutrients*. 2013;5(7):2314-2332. doi:10.3390/nu5072314.
5. Do T, Anh, Hoang L, Shihavong K, Uy N, quang huy N. In vitro Antibacterial Activity of Quercetin Containing Extract from Hibiscus Sabdariffa L. Calyxes. *VNU Journal of Science: Natural Sciences and Technology*. 2016;32(1S):147-152.
6. Nguyen KNH, Nguyen N-VT, Kim KH. Determination of phenolic acids and flavonoids in leaves, calyces, and fruits of *Physalis angulata* L. in Viet Nam. *Pharmacia*. 2021; 68(2): 501-509. doi: 10.3897/pharmacia.68.e66044.
7. Chac L, Thinh B, Huong H, Kiet N, Huan L. Quantification of quercetin, isorhamnetin and ferulic acid in dry extract of *Anoectochilus setaceus* Blume from Vietnam. *International Journal of Botany Studies*. 2019;4(5):14-18.
8. Nguyen D, Herent M-F, Le T-B, Hue B, Hang B, Thi-Thanh-Huong D, et al. Isolation of Quercetin-3-O-sulfate and quantification of major compounds from *Psidium guajava* L. from Vietnam. *Journal of Food Composition and Analysis*. 2022;115:104928. doi:10.1016/j.jfca.2022.104928.
9. Su Z-Y, Lai B-A, Lin Z-H, Wei G-J, Huang S-H, Tung Y-C, et al. Water extract of lotus leaves has hepatoprotective activity by enhancing Nrf2- and epigenetics-mediated cellular antioxidant capacity in mouse hepatocytes. *Journal of Functional Foods*. 2022;99:105331. doi:10.1016/j.jff.2022.105331.
10. Zhu X, Xiong T, Liu P, Guo X, Xiao L, Zhou F, et al. Quercetin ameliorates HFD-induced NAFLD by promoting hepatic VLDL assembly and lipophagy via the IRE1 $\alpha$ /XBP1s pathway. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*. 2018;114:52-60.
11. Carson C, Lawson HA. Epigenetics of metabolic syndrome. *Physiological genomics*. 2018;50(11):947-955.
12. Aguilar-Zarate P, Aguirre Joya JA, Govea Salas M, Rodriguez R. Nutrigenetics and nutrigenomics: a general review. 2014. p. 383-401.
13. Dabeek WM, Marra MV. Dietary Quercetin and Kaempferol: Bioavailability and Potential Cardiovascular-Related Bioactivity in Humans. *Nutrients*. 2019;11(10). doi:10.3390/nu11102288.
14. Nile A, Nile SH, Shin J, Park G, Oh JW. Quercetin-3-Glucoside Extracted from Apple Pomace Induces Cell Cycle Arrest and Apoptosis by Increasing Intracellular ROS Levels. *International journal of molecular sciences*. 2021;22(19). doi:10.3390/ijms221910749.
15. Larson A, Symons J, Jalili T. Therapeutic Potential of Quercetin to Decrease Blood Pressure: Review of Efficacy and Mechanisms. *Advances in nutrition (Bethesda, Md)*. 2012;3:39-46.
16. Sezer ED, Oktay LM, Karadaş E, Memmedov H, Selvi Gunel N, Sözmen E. Assessing Anticancer Potential of Blueberry Flavonoids, Quercetin, Kaempferol, and Gentsic Acid, Through Oxidative Stress and Apoptosis Parameters on HCT-116 Cells. *Journal of medicinal food*. 2019;22(11):1118-26. doi:10.1089/jmf.2019.0098.
17. Haytowitz. DB, Wu. X, Bhagwat. S. USDA Database for the Flavonoid Content of Selected Foods (Release 3.3): U.S. Department of Agriculture. [Online]. 2018. Available from: <https://www.ars.usda.gov/northeast->

- area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/methods-and-application-of-food-composition-laboratory/mafcl-site-pages/flavonoids/.
18. Yasin G, Jasim S, Mahmudiono T, Al-Shawi S, Shichiyakh R, Shoukat S, et al. Investigating the effect of garlic (*Allium sativum*) essential oil on foodborne pathogenic microorganisms. *Food Science and Technology*. 2022;42. doi:10.1590/fst.03822.
19. Yang J, Chen X, Rao S, Li Y, Zang Y, Zhu B. Identification and Quantification of Flavonoids in Okra (*Abelmoschus esculentus* L. Moench) and Antiproliferative Activity In Vitro of Four Main Components Identified. *Metabolites*. 2022;12(6). doi:10.3390/metabo12060483.
20. Hwang IK, Lee CH, Yoo KY, Choi JH, Park OK, Lim SS, et al. Neuroprotective effects of onion extract and quercetin against ischemic neuronal damage in the gerbil hippocampus. *Journal of medicinal food*. 2009;12(5):990-5. doi:10.1089/jmf.2008.1400.
21. Verma RJ, Sangai NP. The ameliorative effect of black tea extract and quercetin on bisphenol A-induced cytotoxicity. *Acta poloniae pharmaceutica*. 2009;66(1):41-4.
22. Soleas GJ, Grass L, Josephy PD, Goldberg DM, Diamandis EP. A comparison of the anticarcinogenic properties of four red wine polyphenols. *Clinical Biochemistry*. 2002;35(2):119-24. doi:10.1016/S0009-9120(02)00275-8.
23. 정창호, 정희록, 최성길 and 허호진. Antioxidant and neuronal cell protective effects of aqueous extracts from lotus leaf tea. *농업생명과학연구*. 2012;46(2):115-127.
24. Naeem mm. Health Benefits of Pomegranate (Peel & Juice) and Preparation of Functional Pomegranate Drink Using Probiotic *Lactobacillus Plantarum*. *J Food Technology Research Journal*. 2023;1(2):77-91.
25. Betanzos-Cabrera G, Montes-Rubio PY, Fabela-Illescas HE, Belefant-Miller H, Cancino-Diaz JC. Antibacterial activity of fresh pomegranate juice against clinical strains of *Staphylococcus epidermidis*. *Food & nutrition research*. 2015;59:27620. doi:10.3402/fnr.v59.27620.
26. Chavez-Santoscoy RA, Gutierrez-Urbe JA, Serna-Saldivar SO. Phenolic Composition, Antioxidant Capacity and In Vitro Cancer Cell Cytotoxicity of Nine Prickly Pear (*Opuntia* spp.) Juices. *Plant Foods for Human Nutrition*. 2009;64(2):146-152.
27. Sambandam B, Devasena T, Arivarasan A, Raman P. Extraction and isolation of flavonoid quercetin from the leaves of *Trigonella foenum-graecum* and their anti-oxidant activity. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2016;8:120-124.
28. Weth FR, Hoggarth GB, Weth AF, Paterson E, White MPJ, Tan ST, et al. Unlocking hidden potential: advancements, approaches, and obstacles in repurposing drugs for cancer therapy. *British Journal of Cancer*. 2024;130(5):703-715.
29. Salehi B, Machin L, Monzote L, Sharifi-Rad J, Ezzat SM, Salem MA, et al. Therapeutic Potential of Quercetin: New Insights and Perspectives for Human Health. *ACS omega*. 2020;5(20):11849-11872. doi:10.1021/acsomega.0c01818.
30. Arnold M, Abnet CC, Neale RE, Vignat J, Giovannucci EL, McGlynn KA, et al. Global Burden of 5 Major Types of Gastrointestinal Cancer. *Gastroenterology*. 2020;159(1):335-49.e15. doi:10.1053/j.gastro.2020.02.068.
31. Syn G, Blackwell JM, Jamieson SE. Chapter 19 - Epigenetics in Infectious Diseases. In: García-Giménez JL, editor. *Epigenetic Biomarkers and Diagnostics*. Boston: Academic Press; 2016. p. 377-400.
32. Stanland LJ, Luftig MA. The Role of EBV-Induced Hypermethylation in Gastric Cancer Tumorigenesis. *Viruses*. 2020;12(11). doi:10.3390/v12111222.
33. Zhang BG, Hu L, Zang MD, Wang HX, Zhao W, Li JF, et al. *Helicobacter pylori* CagA induces tumor suppressor gene hypermethylation by upregulating DNMT1 via AKT-NFκB pathway in gastric cancer development. *Oncotarget*. 2016;7(9):9788-9800. doi:10.18632/oncotarget.7125.
34. Ferro A, Morais S, Pelucchi C, Aragonés N, Kogevinas M, López-Carrillo L, et al. Smoking and *Helicobacter pylori* infection: an individual participant pooled analysis (Stomach Cancer Pooling- StoP Project). *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation (ECP)*. 2019;28(5):390-396. doi:10.1097/cej.0000000000000471.

35. Camargo MC, Koriyama C, Matsuo K, Kim WH, Herrera-Goepfert R, Liao LM, et al. Case-case comparison of smoking and alcohol risk associations with Epstein-Barr virus-positive gastric cancer. *International journal of cancer*. 2014;134(4):948-953. doi:10.1002/ijc.28402.
36. Liu Y, Li Y, Jiang Y, Zheng X, Wang T, Li J, et al. Quercetin Promotes Apoptosis of Gastric Cancer Cells through the EGFR-ERK Signaling Pathway. *Journal of Food Biochemistry*. 2024;2024(1):9945178. doi:10.1155/2024/9945178.
37. Łukasiewicz S, Czezelewski M, Forma A, Baj J, Sitarz R, Stanisławek A. Breast Cancer-Epidemiology, Risk Factors, Classification, Prognostic Markers, and Current Treatment Strategies-An Updated Review. *Cancers*. 2021;13(17). doi:10.3390/cancers13174287.
38. Kashyap D, Pal D, Sharma R, Garg VK, Goel N, Koundal D, et al. Global Increase in Breast Cancer Incidence: Risk Factors and Preventive Measures. *BioMed research international*. 2022;2022:9605439. doi:10.1155/2022/9605439.
39. Lukong KE. Understanding breast cancer - The long and winding road. *BBA clinical*. 2017;7:64-77. doi:10.1016/j.bbacli.2017.01.001.
40. Le Huyen Ai T, Lao Duc T, Truong Kim P. DNA Hypermethylation in Breast Cancer. In: Phuc Van P, editor. Breast cancer (Tokyo, Japan). Rijeka: IntechOpen; 2017. p. Ch. 7.
41. Chen SH, Huang WT, Kao WC, Hsiao SY, Pan HY, Fang CW, et al. O6-methylguanine-DNA methyltransferase modulates cisplatin-induced DNA double-strand breaks by targeting the homologous recombination pathway in nasopharyngeal carcinoma. *Journal of biomedical science*. 2021;28(1):2. doi:10.1186/s12929-020-00699-y.
42. Uramova S, Kubatka P, Dankova Z, Kapinova A, Zolakova B, Samec M, et al. Plant natural modulators in breast cancer prevention: status quo and future perspectives reinforced by predictive, preventive, and personalized medical approach. *EPMA Journal*. 2018;9(4):403-419. doi:10.1007/s13167-018-0154-6.
43. Pirouzpanah S, Taleban FA, Mehdipour P, Atri M. Association of folate and other one-carbon related nutrients with hypermethylation status and expression of RARB, BRCA1, and RASSF1A genes in breast cancer patients. *Journal of molecular medicine (Berlin, Germany)*. 2015;93(8):917-934. doi:10.1007/s00109-015-1268-0.
44. Kundur S, Prayag A, Selvakumar P, Nguyen H, McKee L, Cruz C, et al. Synergistic anticancer action of quercetin and curcumin against triple-negative breast cancer cell lines. *J Cell Physiol*. 2019;234(7):11103-11118. doi:10.1002/jcp.27761.
45. Nguyen L, Lee Y-H, Sharma A, Park J, Jagga S, Sharma G, et al. Quercetin induces apoptosis and cell cycle arrest in triple-negative breast cancer cells through modulation of Foxo3a activity. *Korean Journal of Physiology and Pharmacology*. 2017;21:205-213. doi:10.4196/kjpp.2017.21.2.205.
46. Luo T, Liang Y, Chen L, Sun H, Chen Q. Effect of quercetin on the progression of breast cancer in mice with chronic stress by regulating the polarization of microglia. *Journal of Functional Foods*. 2024; 119:106294. doi:10.1016/j.jff.2024.106294.
47. Prieto-Vila M, Shimomura I, Kogure A, Usuba W, Takahashi RU, Ochiya T, et al. Quercetin Inhibits Lef1 and Resensitizes Docetaxel-Resistant Breast Cancer Cells. *Molecules (Basel, Switzerland)*. 2020;25(11). doi:10.3390/molecules25112576.
48. Dammann RH, Richter AM, Jiménez AP, Woods M, Küster M, Witharana C. Impact of Natural Compounds on DNA Methylation Levels of the Tumor Suppressor Gene RASSF1A in Cancer. *International journal of molecular sciences*. 2017;18(10). doi:10.3390/ijms18102160.
49. Saloni, Goel H, Rawat K, Syeda S, Shrivastava A, Jha A. Demethylation of RASSF1A Gene by Quercetin and Eugenol in HeLa Cancer Cell Line. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2019:29.
50. Shang Z, Kouznetsova VL, Tsigelny IFJJoI. Human Papillomavirus (HPV) Viral Proteins Substitute for the Impact of Somatic Mutations by Affecting Cancer-Related Genes: Meta-analysis and Perspectives. 2020. doi: 10.29245/2689-9981/2020/1.1157.
51. Yuan CH, Filippova M, Tungteakkhun SS, Duerksen-Hughes PJ, Krstenansky JL. Small molecule inhibitors of the HPV16-E6 interaction with caspase 8. *Bioorganic &*

- medicinal chemistry letters*. 2012;22(5):2125-2129. doi:10.1016/j.bmcl.2011.12.145.
52. Murata M, Komatsu S, Miyamoto E, Oka C, Lin I, Kumazoe M, et al. Quercetin up-regulates the expression of tumor-suppressive microRNAs in human cervical cancer. *Bioscience of microbiota, food and health*. 2023;42(1):87-93. doi:10.12938/bmfh.2022-056.
53. Mengstie MA, Chekol Abebe E, Behaile Teklemariam A, Tilahun Mulu A, Agidew MM, Teshome Azezew M, et al. Endogenous advanced glycation end products in the pathogenesis of chronic diabetic complications. *Front Mol Biosci*. 2022;9:1002710. doi:10.3389/fmolb.2022.1002710.
54. Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, et al. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Research and Clinical Practice*. 2022;183:109119. doi:10.1016/j.diabres.2021.109119.
55. Dhanya R, Arya AD, Nisha P, Jayamurthy P. Quercetin, a Lead Compound against Type 2 Diabetes Ameliorates Glucose Uptake via AMPK Pathway in Skeletal Muscle Cell Line. *Frontiers in pharmacology*. 2017;8:336. doi:10.3389/fphar.2017.00336.
56. Niu Y, Wang T, Liu S, Yuan H, Li H, Fu L. Exercise-induced GLUT4 transcription via inactivation of HDAC4/5 in mouse skeletal muscle in an AMPK $\alpha$ 2-dependent manner. *Biochimica et biophysica acta Molecular basis of disease*. 2017;1863(9):2372-2381. doi:10.1016/j.bbadis.2017.07.001.
57. Zhang L, Wang X, Chang L, Ren Y, Sui M, Fu Y, et al. Quercetin improves diabetic kidney disease by inhibiting ferroptosis and regulating the Nrf2 in streptozotocin-induced diabetic rats. *Renal failure*. 2024;46(1):2327495. doi:10.1080/0886022x.2024.2327495.
58. Li Y, Wang L, zhang M, Huang K, Yao Z, Rao P, et al. Advanced glycation end products inhibit the osteogenic differentiation potential of adipose-derived stem cells by modulating Wnt/ $\beta$ -catenin signalling pathway via DNA methylation. *Cell Proliferation*. 2020;52(2):e12540. doi:10.1111/cpr.12834.
59. Zhang Z, Yang X, Song YQ, Tu J. Autophagy in Alzheimer's disease pathogenesis: Therapeutic potential and future perspectives. *Ageing research reviews*. 2021;72:101464. doi:10.1016/j.arr.2021.101464.
60. Pant R, Kabeer SW, Sharma S, Kumar V, Patra D, Pal D, et al. Pharmacological inhibition of DNMT1 restores macrophage autophagy and M2 polarization in Western diet-induced nonalcoholic fatty liver disease. *J Biol Chem*. 2023;299(6):104779. doi:10.1016/j.jbc.2023.104779.
61. Lei P, Hu Y, Gao P, Ding Q, Yan J, Zhao J, et al. Sulforaphane Ameliorates Hepatic Lipid Metabolism via Modulating Lipophagy In Vivo and In Vitro. *Journal of agricultural and food chemistry*. 2022;70(48):15126-15133. doi:10.1021/acs.jafc.2c06311.
62. Cao H, Jia Q, Yan L, Chen C, Xing S, Shen D. Quercetin Suppresses the Progression of Atherosclerosis by Regulating MST1-Mediated Autophagy in ox-LDL-Induced RAW264.7 Macrophage Foam Cells. *International journal of molecular sciences*. 2019;20(23). doi:10.3390/ijms20236093.
63. Chang DPS, Guan XL. Metabolic Versatility of Mycobacterium tuberculosis during Infection and Dormancy. *Metabolites*. 2021;11(2). doi:10.3390/metabo11020088.
64. Sengupta S, Nayak B, Meuli M, Sander P, Mishra S, Sonawane A. Mycobacterium tuberculosis Phosphoribosyltransferase Promotes Bacterial Survival in Macrophages by Inducing Histone Hypermethylation in Autophagy-Related Genes. *Frontiers in cellular and infection microbiology*. 2021;11:676456. doi:10.3389/fcimb.2021.676456.
65. Shukla H, Kumar V, Singh AK, Rastogi S, Khan SR, Siddiqi MI, et al. Isocitrate lyase of Mycobacterium tuberculosis is inhibited by quercetin through binding at N-terminus. *International journal of biological macromolecules*. 2015;78:137-141. doi:10.1016/j.ijbiomac.2015.04.005.
66. Zhu X, Li D, Du Y, He W, Lu Y. DNA hypermethylation-mediated downregulation of antioxidant genes contributes to the early onset of cataracts in highly myopic eyes. *Redox biology*. 2018;19:179-189. doi:10.1016/j.redox.2018.08.012.
67. Francisco BM, Salvador M, Amparo N. Oxidative stress in myopia. *Oxidative medicine and cellular longevity*.

- 2015;2015:750637.  
doi:10.1155/2015/750637.
68. Pawlowska E, Szczepanska J, Koskela A, Kaarniranta K, Blasiak J. Dietary Polyphenols in Age-Related Macular Degeneration: Protection against Oxidative Stress and Beyond. *Oxidative medicine and cellular longevity*. 2019;2019:9682318. doi:10.1155/2019/9682318.
69. Davis CK, Vemuganti R. Antioxidant therapies in traumatic brain injury. *Neurochemistry international*. 2022;152:105255. doi:10.1016/j.neuint.2021.105255.
70. Chen X, Chen C, Fan S, Wu S, Yang F, Fang Z, et al. Omega-3 polyunsaturated fatty acid attenuates the inflammatory response by modulating microglia polarization through SIRT1-mediated deacetylation of the HMGB1/NF- $\kappa$ B pathway following experimental traumatic brain injury. *J Neuroinflammation*. 2018;15(1):116. doi:10.1186/s12974-018-1151-3.
71. Chen X, Wu S, Chen C, Xie B, Fang Z, Hu W, et al. Omega-3 polyunsaturated fatty acid supplementation attenuates microglial-induced inflammation by inhibiting the HMGB1/TLR4/NF- $\kappa$ B pathway following experimental traumatic brain injury. *J Neuroinflammation*. 2017;14(1):143. doi:10.1186/s12974-017-0917-3.
72. Wang Z, Guo W, Yi F, Zhou T, Li X, Feng Y, et al. The Regulatory Effect of SIRT1 on Extracellular Microenvironment Remodeling. *International journal of biological sciences*. 2021;17(1):89-96. doi:10.7150/ijbs.52619.
73. Yang T, Kong B, Gu JW, Kuang YQ, Cheng L, Yang WT, et al. Anti-apoptotic and anti-oxidative roles of quercetin after traumatic brain injury. *Cellular and molecular neurobiology*. 2014;34(6):797-804. doi:10.1007/s10571-014-0070-9.
74. Kom HH, Nageshwar D, Srilatha K, Reddy K. Protective effect of quercetin on weight drop injury model-induced neuroinflammation alterations in brain of mice. *Journal of Applied Pharmaceutical Science*. 2019;9:96-103. doi:10.7324/JAPS.2019.90412.
75. Wang Y, Wang H. The emerging role of histone deacetylase 1 in allergic diseases. *Frontiers in immunology*. 2022;13:1027403. doi:10.3389/fimmu.2022.1027403.
76. Ke X, Chen Z, Wang X, Kang H, Hong S. Quercetin improves the imbalance of Th1/Th2 cells and Treg/Th17 cells to attenuate allergic rhinitis. *Autoimmunity*. 2023;56(1):2189133. doi:10.1080/08916934.2023.2189133.
77. Bar-El Dadon S, Reifen R. Chapter 55 - Epigenetics in Food Allergies: The Missing Piece of the Puzzle. In: Caterina RDE, Martinez JA, Kohlmeier M, editors. *Principles of Nutrigenetics and Nutrigenomics*: Academic Press; 2020. p. 403-409.