

## NUTRITIONAL ROLE OF RICE BRAN IN CARDIOVASCULAR HEALTH

Lam Khac Ky✉, Vien Vu Bao An, Vo Tan Dat

*Industrial University of Ho Chi Minh City, Vietnam*

### ABSTRACT

Cardiovascular diseases are a major cause of death globally, including in Vietnam. In recent years, there has been a rise in younger people developing cardiovascular diseases due to unhealthy lifestyles and diets. The rice bran, a byproduct of rice milling, was traditionally used as animal feed. However, recent research has shown that rice bran contains valuable nutrients such as lipids (15-22%), carbohydrates (34.1-52.2%), fiber (7-14%), and protein (10-16%), as well as beneficial bioactive compounds like flavonoids, ferulic acid, policosanol, and feruloylated oligosaccharides. These components have demonstrated potential in preventing and managing cardiovascular diseases. This review presents the nutritional components of rice bran, the rice bran products available on the market, and research on how rice bran affects cardiovascular health. Based on this research, rice bran could be incorporated into food products to help reduce the risk of cardiovascular diseases.

**Keywords:** *rice bran, chemical composition, cardiovascular diseases.*

-----

### I. INTRODUCTION

Cardiovascular diseases (CVDs) are a group of disorders of the heart and blood vessels. They include coronary heart disease, cerebrovascular disease, peripheral arterial disease, rheumatic heart disease, congenital heart disease, and deep vein thrombosis and pulmonary embolism [1]. The World Health Organization reported that in 2019, CVDs accounted for 39.5% of deaths globally. In Vietnam, data from the Ministry of Health indicates a rising number of CVD cases over the years, with mortality rates increasing from 127.3 per 100,000 people in 2000 to 164.9 per 100,000 people in 2022 [2]. Cardiovascular diseases (CVDs) are the leading cause of death in Vietnam, accounting for approximately 200,000

deaths annually, or 33% of all fatalities (Vietnam Ministry of Health, 2023). These alarming statistics highlight the urgent need for effective interventions and public health strategies to prevent and manage CVDs in Vietnam [57]. Presently, cardiovascular diseases are not only the leading cause of death but also increasingly affect younger individuals, with myocardial infarctions occurring more frequently in people in their 30s and hypertension affecting 16.5% of those under 40 [3]. The rising incidence of cardiovascular diseases among younger populations can be attributed to unhealthy habits, lifestyles, and diets such as high consumption of animal fats, processed foods, and stimulants like alcohol and tobacco.

✉ Corresponding author: Lam Khac Ky  
Email: lamkhacky@iuh.edu.vn  
Doi:10.56283/1859-0381/781

Submitted: August 30, 2024  
Revised: December 2, 2025  
Accepted: December 9, 2025  
Online first: December 9, 2025

High-risk individuals often include those who are overweight or obese, have large waistlines, have dyslipidemia, and particularly smokers [3].

Preventing cardiovascular disease (CVD) requires addressing major behavioral risk factors. The World Health Organization (WHO) recommends the following measures:

**Avoid tobacco use:** smoking is a major risk factor for CVD. Quitting smoking and avoiding exposure to second-hand smoke can significantly reduce the risk [58].

**Maintain a healthy diet:** consuming plenty of fruits, vegetables, whole grains, and nuts while limiting saturated fats, trans fats, salt, and sugar can help prevent CVD. WHO recommends eating at least 400 grams of fruits and vegetables per day and reducing salt intake to less than 5 grams per day [58].

**Engage in regular physical activity:** participating in at least 150 minutes of moderate-intensity physical activity per week is recommended to maintain cardiovascular health [58].

Rice bran (RB) is a type of cereal flour produced as a by-product of rice milling. It is rich in nutrients, containing 15-22% lipids, 34.1-52.2% carbohydrates, 7-14%

fiber, 6.6-9.9% ash, and 10-16% protein [4]. Additionally, rice bran contains both soluble and insoluble fibers, vitamins especially B vitamins and minerals [5]. The primary fats in rice bran are short-chain and unsaturated. Furthermore, its protein comprises nearly all essential amino acids required by the body, such as lysine, histidine, threonine, cystine, valine, methionine, isoleucine, leucine, and phenylalanine [6].

Recently, rice bran has garnered interest for its numerous health benefits. Studies have shown rice bran's benefits include protecting and preventing intestinal immune system disorders, improving bowel dysfunction, lowering blood lipid levels, controlling blood glucose levels, and guarding against certain cancers and cardiovascular diseases. Beyond improving bowel dysfunction, rice bran's role in disease prevention and protection is of significant research interest. Multiple studies indicate that rice bran can reduce the risk of cardiovascular disease. This article aims to summarize the role of rice bran in cardiovascular diseases and elucidate the components in rice bran that contribute to reducing cardiovascular disease risks.

## II. METHODS

The selected documents were research articles published in reputable scientific journals in

<https://pubmed.ncbi.nlm.nih.gov> and

<https://hinari.summon.serialssolutions.com>.

The selected documents included full research articles, short communications,

and review articles, encompassing the results of experimental studies, clinical trials, and survey data.

Keywords used for the literature search were: rice bran, rice bran composition, rice bran protein, rice bran oil, rice bran and cardiovascular disease, rice bran and heart health

### III. NUTRITIONAL ROLE OF RICE BRAN IN CARDIO-VASCULAR HEALTH

#### 3.1. Rice bran protein

Rice bran protein, although constituting 10-16% of rice bran, has demonstrated significant nutritional and functional properties comparable to proteins in other grains. The protein fraction in rice bran includes albumin (37-40%), globulin (21-36%), prolamin (3-5%), and glutelin (22-36%) [6, 7].

**Antihypertensive Effects:** rice bran protein and its hydrolysates have been found to exhibit antihypertensive properties. Peptides such as Tyr-Tyr and Tyr-Ser-Lys are reported to inhibit angiotensin-converting enzyme (ACE), contributing to blood pressure reduction [10, 55]. Thermolysin-digested rice bran (TRB), Protease G6-digested rice bran, and trypsin-digested rice bran have also been shown to lower systolic blood pressure (SBP) in experimental models [51, 55]. Additionally, a clinical trial demonstrated that rice bran containing the novel peptide Leu-Arg-Ala significantly reduced SBP in

hypertensive individuals without adverse effects [10].

**Antidiabetic Effects:** peptides derived from rice bran protein can inhibit amylase,  $\alpha$ -glucosidase,  $\beta$ -glucosidase, and dipeptidyl peptidase-IV (DPP-IV), suggesting potential benefits in managing blood glucose levels [52].

**Cognitive Improvement:** rice-memolin, a pentapeptide derived from rice bran protein, has shown cognitive enhancement in animal studies. It promotes neurogenesis and the expression of neurotrophic factors in the hippocampus, which may aid in mitigating cognitive decline [51].

**Antimicrobial Properties:** peptides from rice bran protein exhibit antimicrobial properties, potentially acting against various bacterial strains [52].

**Enhancement of Iron Availability:** peptides derived from rice bran protein enhance iron bioavailability, supporting improved nutritional outcomes [53].

#### 3.2. Rice bran oil

For many years, health organizations have advocated and encouraged people to reduce their intake of saturated fat to lower the risk of coronary heart disease [14, 15]. The reason is that saturated fats can raise blood cholesterol levels more than polyunsaturated fats [16]. Rice bran oil (RBO) contains a high amount of monounsaturated fatty acids (42%) and polyunsaturated fatty acids (40%; 38% omega-6 fatty acids and 2% omega-3

fatty acids), with a low level of saturated fat (18%) [17]. Studies have reported that dietary supplementation with RBO can reduce total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and triglycerides (TG) while increasing high-density lipoprotein cholesterol (HDL-C) levels in the blood. Therefore, commercial RBO products offer potential health benefits for managing hypercholesterolemia [18]. Research

indicates that incorporating rice bran oil into daily diets reduces serum cholesterol levels and LDL cholesterol fractions. Since a 1% decrease in cholesterol can lead to a 2% reduction in the risk of heart attack, rice bran oil may be beneficial for patients with heart disease [19]. Additionally, the ratio of SFA:MUFA:PUFA in rice bran oil is close to the recommended 10:15:10 ratio by the American Heart Association for patients with cardiovascular conditions to prevent hypertension, atherosclerosis, and high cholesterol [20, 21]. Dietary adjustment is the most effective first step to address hyperlipidemia and atherosclerosis. The development of atherosclerotic lesions is significantly prevented if the diet includes a mixture of phytochemicals that lower cholesterol and antioxidants [22].

Today, the use of rice bran oil as a cooking oil has become increasingly popular among consumers due to its anti-inflammatory, anti-dyslipidemic, and anti-thrombotic properties (Table 1). Rao and SUGASini [35] have demonstrated that the unsaponifiable fraction of rice bran oil significantly contributes to its anti-inflammatory activity. Inflammatory mediators such as ROS (O<sub>2</sub> and NO<sub>2</sub>), eicosanoids (PGE<sub>2</sub>, TXB<sub>2</sub>, LTB<sub>4</sub>, and LTC<sub>4</sub>), cytokines (TNF- $\alpha$  and IL-6), and hydrolytic enzymes (collagenase, elastase, and hyaluronidase) were significantly reduced in isolated peritoneal macrophages from male Wistar rats fed crude rice bran oil containing saponifiable fractions [35]. Francisqueti and Minatel [35] investigated the therapeutic effects of  $\gamma$ -oryzanol derived from rice bran oil in preventing cardio-renal metabolic

syndrome. Wistar rats fed a high-fat diet for 20 weeks, with or without  $\gamma$ -oryzanol supplementation, showed that the  $\gamma$ -oryzanol-supplemented diet prevented weight gain, increased blood triglycerides, and improved systolic and diastolic function [36].

In humans, testing the effects of supplementing diets with palm oil, rice bran oil, and coconut oil in subjects with normal blood lipid levels revealed an increased susceptibility to excessive chylomicron development (small fat particles that transport lipids to tissues) in response to meals rich in saturated fatty acids from palm oil, thereby raising the risk of atherosclerosis [37]. However, those consuming rice bran oil or coconut oil did not develop atherosclerosis risk factors [37]. In hyperlipidemic patients, a diet supplemented with rice bran oil has been shown to reduce body weight, body mass index, waist circumference, and hip circumference [38], as well as total cholesterol, LDL cholesterol, and the atherogenic index of total cholesterol/HDL cholesterol [38]. Kuriyan and Gopinath [39] also confirmed that rice bran oil significantly reduced total cholesterol and triglycerides in the plasma of hyperglycemic patients compared to common cooking oils like sunflower oil. Additionally, mixing rice bran oil with other cooking oils such as sesame oil [40, 41] or palm oil [42] also shows so

From the information above, it is evident that rice bran contains components that can intervene and protect cardiovascular health. Further studies need to be done *in vitro* and *in vivo* studies on the bioactive compounds of rice bran.

**Table 1.** Cardioprotective roles of *in vitro* and *in vivo* bioactive compounds derived from rice bran.

Compound/Source	Methods	Key Findings
Rice Bran Enzyme Extract [43–45, 47]	<i>In vivo</i> animal studies (ApoE <sup>−/−</sup> mice, Zucker rats, Wistar rats)	- Reduced serum lipids, atherosclerotic plaque development, hepatic steatosis, and inflammation - Improved endothelial function and increased antioxidant activity
Feruloylated Oligosaccharides [31]	<i>In vitro</i> culture (RAW264.7 cells)	- Increased IL-10; decreased TNF- $\alpha$ , IL-1 $\beta$ , IL-6, NO, and PGE2
Rice Bran Policosanol [32]	<i>In vivo</i> (Sprague-Dawley rats)	- Reduced platelet adhesion and cellular protein secretion
Resveratrol and Rice Bran Blend [46]	<i>In vivo</i> (Sprague-Dawley rats)	- Reduced infarct size, increased survival signaling, and enhanced anti-apoptotic protein activity
Bioactive Rice Bran Peptides [49]	<i>In vitro</i> (HUVECs)	- Reduced H <sub>2</sub> O <sub>2</sub> -induced cell death and oxidative stress
Hydrolyzed Rice Bran Protein [50–51]	<i>In vivo</i> (Sprague-Dawley rats, renovascular hypertension model)	- Improved endothelial function, reduced hypertension, hyperglycemia, and insulin resistance
Rice Bran Oil [52–54]	<i>In vivo</i> (Wistar rats, human studies)	- Reduced inflammatory mediators, improved lipid profiles, and decreased atherosclerosis risk - Lowered LDL, triglycerides, and C-reactive protein levels
Rice Bran Blends [40–42]	<i>In vivo</i> (human studies, hypertensive and hyperlipidemic participants)	- Lowered LDL, total cholesterol, triglycerides; improved HDL levels

### 3.3. Biologically active compounds

Atherosclerosis is one of the most significant conditions in cardiovascular disease (CVD), characterized by the accumulation of plaques on arterial walls. This leads to the hardening and narrowing of arteries, impairing blood circulation [23, 24]. The underlying

cause of atherosclerosis involves the dysfunction of endothelial cells that line the inner surface of blood vessels [25], which is triggered by free radicals, adhesive molecules (reactive nitrogen species, reactive oxygen species), inflammatory factors, and thrombosis-

promoting agents [26]. Additionally, the interaction between the endothelium, leukocytes, and inflammatory mediators such as ICAM-1, VCAM-1, IL-1 $\beta$ , and IL-6 can contribute to the development of atherosclerosis [27, 28, 29].

Researchers have identified several bioactive compounds in rice bran that can reduce thrombus formation and vascular inflammation through their antioxidant activities. These include acylated stryly glucoside [30], flavonoids [31], ferulic acid [32], and policosanols [33], as well as components in plasma like leukocytes and platelets [34]. By scavenging free radicals and exhibiting antioxidant properties, these bioactive compounds in rice bran help reduce and prevent the onset of cardiovascular disease.

In a study on rice metabolism, researchers found bioactive flavonoids and their isomers, such as tricin and tricin 7-O-rutinoside, as well as tricin 7-O- $\beta$ -D-glucopyranoside, predominantly concentrated in rice leaves and bran [31]. Additionally, rice bran contains tricin 4'-O-(erythron- $\beta$ -guaiacylglyceryl) ether

and isoscoparin 2''-O-(-(E)-feruloyl) glucopyranoside, which can inhibit excessive nitric oxide (NO) production by macrophages and scavenge the 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical [32]. Feruloylated oligosaccharides (conjugates of ferulic acid) extracted from rice bran have also been recognized for their ability to inhibit atherosclerotic agents such as TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and NO, and to stimulate IL-10 production in lipopolysaccharide (LPS)-stimulated macrophages (RAW264.7) in a dose-dependent manner (0.1–100  $\mu$ g/mL) [32]. Furthermore, in the context of vascular inflammation, the production of prostaglandin E2 (PGE2) can be inhibited by feruloylated oligosaccharides from rice bran (100  $\mu$ g/mL) [32]. Policosanols extracted from rice bran wax has demonstrated anti-thrombotic activity [33]. Mice pretreated with varying concentrations (125–1000  $\mu$ g/mL) of policanol extract showed a dose-dependent reduction in platelet aggregation with collagen and secreted cellular proteins [33].

### 3.4 Effects of rice bran and its products on cardiovascular disease in humans

Cardiovascular disease (CVD) is a leading cause of death globally, with dietary interventions playing a critical role in its prevention and management. Rice bran (RB), a byproduct of rice milling, is rich in bioactive compounds and has gained attention for its potential health benefits, particularly in cardiovascular health [38, 39]. Rice bran contains essential nutrients, including lipids, proteins, fibers, and bioactive components such as  $\gamma$ -oryzanol, tocopherols, tocotrienols, and unsaturated fatty acids [55, 56]. These compounds exhibit antioxidant, anti-

inflammatory, and cholesterol-lowering properties, making RB a promising functional food for cardiovascular disease management [36, 56].

**Lipid Profile Improvement:** research indicates that rice bran oil (RBO) can significantly reduce total cholesterol, LDL-C, and triglycerides while increasing HDL-C. Zavoshy et al. (2012) demonstrated that dietary supplementation with RBO improved lipid profiles in hyperlipidemic patients, reducing the risk of atherosclerosis and coronary artery disease [38]. Similarly, Kuriyan et al. (2005) found that daily

consumption of RBO for eight weeks led to substantial improvements in lipid profiles among overweight individuals [39].

**Blood Pressure Regulation:** peptides derived from rice bran have been shown to exhibit antihypertensive properties. Ogawa et al. (2019) conducted a randomized controlled trial demonstrating that participants consuming processed RB experienced significant reductions in systolic blood pressure (SBP) compared to those receiving a placebo [55]. This suggests the potential of RB peptides, such as Leu-Arg-Ala, as natural ACE inhibitors [55, 10].

**Anti-inflammatory and Antioxidant effects** Rice bran oil contains  $\gamma$ -oryzanol and other unsaponifiable fractions that reduce inflammatory markers such as TNF- $\alpha$  and IL-6. Rao and Sugasini (2016) showed that these components decreased oxidative stress and inflammation in animal models [56]. Human studies have also highlighted the role of RBO in improving oxidative stress markers, contributing to vascular health [36, 56].

**Prevention of Atherosclerosis:** francisqueti et al. (2023) studied the effects of  $\gamma$ -oryzanol supplementation on endothelial function and LDL oxidation. Their findings suggest that regular

consumption of RBO can prevent atherosclerosis by reducing oxidative damage and improving vascular endothelial health [36].

**Mechanisms of Cardioprotective Effects** **Cholesterol Reduction:**

$\gamma$ -Oryzanol and unsaturated fatty acids inhibit cholesterol absorption and enhance bile acid excretion [38, 36].

**Antioxidant Properties:** Tocopherols and tocotrienols neutralize free radicals, reducing oxidative stress on endothelial cells [36, 56].

**Anti-inflammatory Effects:** RB bioactives suppress pro-inflammatory cytokines and promote an anti-inflammatory state [10, 56].

**Blood Pressure Modulation:** RB peptides act as ACE inhibitors, promoting vasodilation and reducing hypertension [55, 10].

Rice bran and its derivatives, particularly RBO, offer significant potential in improving cardiovascular health. By modulating lipid profiles, reducing oxidative stress, and lowering blood pressure, RB-based interventions represent an effective dietary strategy for CVD prevention and management. Further research is needed to validate these findings through long-term clinical trials and optimize RB utilization in functional foods [10, 36].

## IV. DISCUSSION

The role of rice bran (RB) and its derivatives in cardiovascular health has been increasingly recognized in recent years. Bioactive components of RB, such as  $\gamma$ -oryzanol, tocopherols, tocotrienols, and unsaturated fatty acids, have demonstrated significant

cardioprotective effects through lipid modulation, antioxidant properties, and anti-inflammatory mechanisms [38, 39, 55]. These compounds collectively address critical risk factors for cardiovascular disease (CVD).



Clinical studies have confirmed that RB and its products can effectively reduce LDL-C levels, increase HDL-C levels, lower blood pressure, and mitigate vascular inflammation [38, 55]. For instance, randomized controlled trials have demonstrated the lipid-lowering effects of rice bran oil (RBO) and the antihypertensive effects of RB-derived peptides, particularly in individuals with metabolic syndrome or hyperlipidemia [36, 39].

Despite these promising findings, several challenges and research gaps remain. Most studies to date have focused on short- to medium-term interventions, leaving the long-term effects of RB consumption on CVD outcomes unclear. Additionally, further research is needed to explore the specific mechanisms of action of RB bioactives in diverse populations and to establish optimal dosing regimens. Potential side effects of prolonged RB supplementation also warrant investigation [10, 56].

As one of the world's largest rice producers, Vietnam generates a substantial amount of rice bran as a byproduct of rice milling. This underutilized resource offers a unique opportunity to develop functional food products aimed at improving public health and reducing the burden of CVD [10]. To fully leverage this potential, Vietnam should consider the following strategies:

## VI. CONCLUSION

Rice bran (RB) and its derivatives, including rice bran oil and rice bran protein, have demonstrated significant potential in preventing and managing cardiovascular diseases. The bioactive compounds in RB, such as  $\gamma$ -oryzanol,

Promote Research and Development: Increased investment in scientific research to better understand the health benefits of RB, particularly its effects on local populations, is essential [10, 56].

Enhance Processing Technologies: Developing advanced methods to stabilize RB and extract its bioactive components efficiently can increase its shelf life and nutritional value [36].

Create Consumer Awareness: Public education campaigns highlighting the health benefits of RB and its derivatives can encourage greater consumption and use in everyday diets [10, 55].

Develop Functional Foods: Encouraging the food industry to incorporate RB into commonly consumed products, such as cooking oils, snacks, and beverages, can make RB-based interventions more accessible [10, 36].

The widespread availability of rice bran in Vietnam represents a significant opportunity for advancing public health through dietary interventions. By promoting RB-based innovations, Vietnam can address the growing prevalence of CVD while establishing itself as a leader in the development of health-focused food products. However, this requires coordinated efforts from researchers, policymakers, and industry stakeholders [10].

tocopherols, tocotrienols, and unsaturated fatty acids, play crucial roles in improving lipid profiles, reducing oxidative stress, and mitigating vascular inflammation.



Incorporating RB as a functional food can effectively reduce the risk of cardiovascular diseases and support a heart-healthy diet. However, to fully realize its benefits, further investments in research, development of advanced processing technologies, and public awareness campaigns are essential.

With its abundant rice production, Vietnam has the opportunity to become a leader in developing innovative health-focused food products based on rice bran. This approach could significantly improve public health while addressing the burden of cardiovascular diseases.

## References

1. World Health Organization. Cardiovascular diseases (CVDs). Accessed December 30, 2024 at: [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)).
2. Bộ Y tế. Cổng thông tin điện tử. Xu hướng tử vong do bệnh tim mạch đang ngày càng tăng tại Việt Nam. Truy cập ngày 15/03/2024 từ: [https://moh.gov.vn/tin-tong-hop/-/asset\\_publisher/k206Q9qkZOqn/content/xu-huong-tu-vong-do-benh-tim-mach-ngay-cang-tang-tai-viet-nam](https://moh.gov.vn/tin-tong-hop/-/asset_publisher/k206Q9qkZOqn/content/xu-huong-tu-vong-do-benh-tim-mach-ngay-cang-tang-tai-viet-nam)
3. Bộ Y tế. Cổng thông tin điện tử. Mỗi năm Việt Nam có 200.000 người tử vong do các bệnh tim mạch. Truy cập ngày 15/03/2024 từ: [https://moh.gov.vn/tin-lien-quan/-/asset\\_publisher/vjYyM7O9aWnX/content/moi-nam-viet-nam-co-200-000-nguoi-tu-vong-do-cac-benh-tim-mach?inheritRedirect=false](https://moh.gov.vn/tin-lien-quan/-/asset_publisher/vjYyM7O9aWnX/content/moi-nam-viet-nam-co-200-000-nguoi-tu-vong-do-cac-benh-tim-mach?inheritRedirect=false)
4. Zheng Y, Gao N, Wu J, Yin B. Rice bran protein: Extraction, nutraceutical properties, and potential applications. In: Rice bran and rice bran oil. *AOCS Press*; 2019:271–293.
5. Tổng cục Thống kê. Báo cáo tình hình kinh tế – xã hội quý IV và năm 2021. 2021.
6. Fabian C, Ju YH. A review on rice bran protein: its properties and extraction methods. *Critical Reviews in Food Science and Nutrition*. 2011;51(9):816–827.
7. Saunders RM. Rice bran: Composition and potential food uses. *Food Reviews International*. 1985;1(3):465–495.
8. Islam S, Nagasaka R, Ohara K, et al. Biological abilities of rice bran-derived antioxidant phytochemicals for medical therapy. *Current Topics in Medicinal Chemistry*. 2011;11(14):1847–1853.
9. Hongu N, Kitts DD, Zawistowski J, et al. Pigmented rice bran and plant sterol combination reduces serum lipids in overweight and obese adults. *Journal of the American College of Nutrition*. 2014;33(3):231–238.
10. Ogawa Y, Shobako N, Fukuhara I, et al. Rice bran supplement containing a functional substance, the novel peptide Leu-Arg-Ala, has anti-hypertensive effects: A double-blind, randomized, placebo-controlled study. *Nutrients*. 2019;11(4):726.
11. Zhang H, Wang J, Liu Y, Gong L, Sun B. Rice bran proteins and their hydrolysates modulate cholesterol metabolism in mice on hypercholesterolemic diets. *Food & Function*. 2016;7(6):2747–2753.
12. Wang J, Shimada M, Nagaoka S. Identification of the active protein in rice bran protein having an inhibitory activity of cholesterol micellar solubility. *Bioscience, Biotechnology, and Biochemistry*. 2017;81(6):1216–1219.
13. Ardiansyah, Ohsaki Y, Shirakawa H, et al. Novel effects of a single administration of ferulic acid on the regulation of blood pressure and the hepatic lipid metabolic profile in stroke-prone spontaneously hypertensive rats. *Journal of Agricultural and Food Chemistry*. 2008;56(8):2825–2830.
14. EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA). Scientific opinion on dietary reference values for fats, including saturated fatty acids, polyunsaturated fatty acids, monounsaturated fatty acids, trans fatty acids, and cholesterol. *EFSA Journal*. 2010;8(3):1461.
15. Eckel RH, Jakicic JM, Ard JD, et al. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014;129(25\_suppl\_2):S76–S99.

16. Kris-Etherton PM, Yu S. Individual fatty acid effects on plasma lipids and lipoproteins: Human studies. *The American Journal of Clinical Nutrition*. 1997;65(5):1628–1644.
18. Park HY, Lee KW, Choi HD. Rice bran constituents: Immunomodulatory and therapeutic activities. *Food & Function*. 2017;8(3):935–943.
19. Bollinedi H, Singh AK, Singh N, et al. Genetic and genomic approaches to address rapid rancidity of rice bran. *Critical Reviews in Food Science and Nutrition*. 2021;61(1):75–84.
20. Chou TW, Ma CY, Cheng HH, et al. A rice bran oil diet improves lipid abnormalities and suppress hyperinsulinemic responses in rats with streptozotocin/nicotinamide-induced type 2 diabetes. *J Clin Biochem Nutr*. 2009;45(1):29–36.
21. Cicero AFG, Gaddi A. Rice bran oil and  $\gamma$ -oryzanol in the treatment of hyperlipoproteinaemias and other conditions. *Phytotherapy Research*. 2001;15(4):277–289.
22. Sohail M, Rakha A, Butt MS, et al. Rice bran nutraceuticals: A comprehensive review. *Crit Rev Food Sci Nutr*. 2017;57(17):3771–3780.
23. Santhakumar AB, Bulmer AC, Singh I. A review of the mechanisms and effectiveness of dietary polyphenols in reducing oxidative stress and thrombotic risk. *Journal of Human Nutrition and Dietetics*. 2014;27(1):1–21.
24. Kumar A. Computational model of blood flow in the presence of atherosclerosis. In: 6th World Congress of Biomechanics (WCB 2010). *Springer Berlin Heidelberg*; 2010:1591–1594.
25. Liao JK. Linking endothelial dysfunction with endothelial cell activation. *J Clin Invest*. 2013;123(2):540–541.
26. Sitia S, Tomasoni L, Atzeni F, et al. From endothelial dysfunction to atherosclerosis. *Autoimmunity Reviews*. 2010;9(12):830–834.
27. Chávez-Sánchez L, Chávez-Rueda K, Legorreta-Haquet MV, et al. The activation of CD14, TLR4, and TLR2 by mmLDL induces IL-1 $\beta$ , IL-6, and IL-10 secretion in human monocytes and macrophages. *Lipids in Health and Disease*. 2010;9(1):1–8.
28. Libby P, Lichtman AH, Hansson GK. Immune effector mechanisms implicated in atherosclerosis: From mice to humans. *Immunity*. 2013;38(6):1092–1104.
29. Hassan GS, Merhi Y, Mourad W. CD40 ligand: A neo-inflammatory molecule in vascular diseases. *Immunobiology*. 2012;217(5):521–532.
30. Nhung BT, Tuyen LD, Linh VA, et al. Rice bran extract reduces the risk of atherosclerosis in post-menopausal Vietnamese women. *Journal of Nutritional Science and Vitaminology*. 2016;62(5):295–302.
31. Yang Z, Nakabayashi R, Mori T, et al. Metabolome analysis of *Oryza sativa* (rice) using liquid chromatography-mass spectrometry for characterizing organ specificity of flavonoids with anti-inflammatory and anti-oxidant activity. *Chemical and Pharmaceutical Bulletin*. 2016;64(7):952–956.
32. Fang HY, Chen YK, Chen HH, et al. Immunomodulatory effects of feruloylated oligosaccharides from rice bran. *Food Chemistry*. 2012;134(2):836–840.
33. Wong WT, Ismail M, Imam MU, Zhang YD. Modulation of platelet functions by crude rice (*Oryza sativa*) bran policosanol extract. *BMC Complementary and Alternative Medicine*. 2016;16(1):1–10.
34. Li KJ, Borresen EC, Jenkins-Puccetti N, et al. Navy bean and rice bran intake alters the plasma metabolome of children at risk for cardiovascular disease. *Frontiers in Nutrition*. 2018;4:71.
35. Rao YPC, Sugasini D, Lokesh B. Dietary gamma oryzanol plays a significant role in the anti-inflammatory activity of rice bran oil by decreasing pro-inflammatory mediators secreted by peritoneal macrophages of rats. *Biochemical and Biophysical Research Communications*. 2016;479(5):747–752.
36. Francisqueti FV, Minatel IO, Ferro AJ, et al. Effect of gamma-oryzanol as a therapeutic agent to prevent cardiorenal metabolic syndrome in animals submitted to high sugar-fat diet. *Nutrients*. 2017;9(9):1299.
37. Irawati D, Mamo JC, Slivkoff-Clark KM, et al. Dietary fat and physiological determinants of plasma chylomicron remnant homeostasis in normolipidemic subjects: Insight into atherogenic risk. *British Journal of Nutrition*. 2017;117(3):403–412.
38. Zavoshy R, Noroozi M, Jahanihashemi H. Effect of low calorie diet with rice bran oil on cardiovascular risk factors in hyperlipidemic patients. *J Res Med Sci*. 2012;17(7):626–631.

39. Kuriyan R, Gopinath N, Vaz M, Kurpad AV. Use of rice bran oil in patients with hyperlipidaemia. *Natl Med J India*. 2005;18(6):292–296.
40. Upadya H, Devaraju C, Joshi SR. Anti-inflammatory properties of blended edible oil with synergistic antioxidants. *IJEM*. 2015;19(4):511–518.
41. Malve H, et al. LDL-cholesterol lowering activity of a blend of rice bran oil and safflower oil in patients with hyperlipidemia. *JIMA*. 2010;108(11):785–788.
42. Devarajan S, Singh R, Chatterjee B, et al. A blend of sesame oil and rice bran oil lowers blood pressure and improves the lipid profile in mild-to-moderate hypertensive patients. *J Clin Lipidol*. 2016;10(2):339–349.
43. Perez-Tenero C, Claro C, Parrado J, Herrera MD, et al. Rice bran enzymatic extract reduces atherosclerotic plaque development and steatosis in high-fat fed ApoE<sup>-/-</sup> mice. *Nutrition*. 2017; 37:22–29.
44. Perez-Tenero C, Herrera M, Laufs U, et al. Food supplementation with rice bran enzymatic extract prevents vascular apoptosis and atherogenesis in ApoE<sup>-/-</sup> mice. *Eur J Nutr*. 2017;56(1):225–236.
45. Justo M, Claro C, Vila E, et al. Microvascular disorders in obese Zucker rats are restored by a rice bran diet. *Nutr Metab Cardiovasc Dis*. 2014;24(5):524–531.
46. Justo ML, Candiracci M, Dantas AP, et al. Rice bran enzymatic extract restores endothelial function and vascular contractility in obese rats by reducing vascular inflammation and oxidative stress. *J Nutr Biochem*. 2013;24(9):1453–1461.
47. Mukherjee S, Ray D, Lekli I, et al. Effects of Longevinex (modified resveratrol) on cardioprotection. *Can J Physiol Pharmacol*. 2010;88(10):1017–1025.
48. Perez-Tenero C, Macià A, de Sotomayor MA, et al. Bioavailability of the ferulic acid-derived phenolic compounds of a rice bran enzymatic extract and their activity against superoxide production. *Food Funct*. 2017;8(6):2165–2174.
49. Juan J, Liu G, Willett WC, Hu FB, et al. Whole grain consumption and risk of ischemic stroke: Results from two prospective cohort studies. *Stroke*. 2017;48(12):3203–3209.
50. Liang Y, Lin Q, Huang P, et al. Rice bioactive peptide binding with TLR4 to overcome H<sub>2</sub>O<sub>2</sub>-induced injury in human umbilical vein endothelial cells through NF- $\kappa$ B signaling. *J Agric Food Chem*. 2018;66(2):440–448.
51. Senaphan K, Sangartit W, Pakdeechote P, et al. Rice bran protein hydrolysates reduce arterial stiffening, vascular remodeling, and oxidative stress in rats fed a high-carbohydrate and high-fat diet. *Eur J Nutr*. 2018; 57(6):219–230.
52. Boonla O, Kukongviriyapan U, Pakdeechote P, et al. Peptides derived from Thai rice bran improve endothelial function in hypertensive rats. *Nutrients*. 2015;7(7):5783–5799.
53. Choi JS, Ron MK, Kim TU, et al. Characteristics of esterified rice bran oil converted by enzymatic esterification. *J Environ Biol*. 2015, 36(5), 1247.
54. Reena M, Krishnakantha TBK. Lowering of platelet aggregation and serum eicosanoid levels in rats fed with a diet containing coconut oil blends with rice bran oil or sesame oil. Prostaglandins Leukot. Essent. *Fatty Acids*. 2010;83(3):151–160.
55. Ogawa Y, Shobako N, & Fukuhara I. Anti-Hypertensive Effects of Peptides Derived from Rice Bran Protein. *Nutrients*. 2019;12(10):3060.
56. Rao YPC & Sugasini D. Anti-inflammatory activity of rice bran oil in animal models. *Biochemical and Biophysical Research Communications*. 2016;479(5):747–752.
57. Bộ Y tế. Công thông tin điện tử. Khoảng 200.000 người Việt tử vong vì tim mạch mỗi năm, 8 lời khuyên để không mắc bệnh này. Truy cập ngày 06/01/2025 từ: [https://moh.gov.vn/hoat-dong-cua-dia-phuong/-/asset\\_publisher/gHbla8vOQDuS/content/khoang-200-000-nguoi-viet-tu-vong-vi-tim-mach-moi-nam-8-loi-khuyen-e-khong-mac-benh-nay](https://moh.gov.vn/hoat-dong-cua-dia-phuong/-/asset_publisher/gHbla8vOQDuS/content/khoang-200-000-nguoi-viet-tu-vong-vi-tim-mach-moi-nam-8-loi-khuyen-e-khong-mac-benh-nay)
58. World Health Organization. Cardiovascular diseases (CVD) in Viet Nam. Accessed January 6, 2025 at: <https://www.who.int/vietnam/health-topics/cardiovascular-disease>.